

Status and future prospects of the NEWS-G experiment

P. Knights* on behalf of the NEWS-G collaboration

School of Physics and Astronomy, University of Birmingham, B15 2TT, UK

* p.r.knights@bham.ac.uk



14th International Conference on Identification of Dark Matter
Vienna, Austria, 18-22 July 2022
doi:[10.21468/SciPostPhysProc.12](https://doi.org/10.21468/SciPostPhysProc.12)

Abstract

The NEWS-G collaboration is searching for light dark matter using spherical proportional counters. Access to the mass range from 50 MeV to 10 GeV is enabled by the combination of low energy threshold, light gaseous targets (H, He, Ne), and highly radio-pure detector construction. Several of the recent developments that have paved the way for the operation of a new 140 cm in diameter, spherical proportional counter are presented. Constructed and commissioned at LSM using 4N copper with a $500\ \mu\text{m}$ electroplated inner layer, the detector is now installed in SNOLAB, where it has begun data taking. Building on the electroplating of the current detector, the collaboration construct its next detector directly in the underground laboratory, using ultra-pure copper electroforming. The design and construction of ECUME, a 140 cm in diameter spherical proportional counter fully electroformed underground is discussed, along with the potential to achieve sensitivity reaching the neutrino floor in light Dark Matter searches with a next-generation detector.



Copyright P. Knights.

This work is licensed under the Creative Commons

[Attribution 4.0 International License](https://creativecommons.org/licenses/by/4.0/).

Published by the SciPost Foundation.

Received 04-10-2022

Accepted 28-04-2023

Published 03-07-2023

doi:[10.21468/SciPostPhysProc.12.024](https://doi.org/10.21468/SciPostPhysProc.12.024)



Check for updates

1 Introduction and status

Direct searches for Dark Matter (DM) have been carried out for several decades, primarily focussed in the mass range 10–1000 GeV, with a recent review presented in Ref. [1]. However, theoretical and experimental interest has recently been attracted to the light-DM region, below a few GeV, which remains relatively unexplored [2–4]. Access to this region of parameter space requires either novel detection channels [5] or lower energy thresholds and light targets [1].

The New Experiments With Spheres - Gas (NEWS-G) collaboration is searching for light-DM using an innovative gaseous detector, the spherical proportional counter [6–9], shown in Figure 1. First results with a $\varnothing 60\text{cm}$ prototype detector operated in the Modane Underground Laboratory (LSM), France, were obtained with a $9.7\ \text{kg}\cdot\text{day}$ exposure with a $\text{Ne}:\text{CH}_4$ gas




Figure 1: Schematic and principle of operation of the spherical proportional counter [6].

mixture. This produced the most stringent constraint at the time on the spin-independent DM-nucleon scattering cross-section, $4.4 \times 10^{-37} \text{ cm}^2$, for a 0.5 GeV mass DM candidate [10]. This data have also recently been used to perform a search for Kaluza-Klein axions [11].

NEWS-G have since constructed a $\varnothing 140 \text{ cm}$ detector [12]. Shown in Figure 2, the detector benefits from increased radiopurity material construction, namely 99.99% pure copper with an ultra-pure copper electroformed internal layer [13], and an improved, compact shielding.



Figure 2: Left: The 140 cm in diameter NEWS-G spherical proportional counter being installed in SNOLAB. Right: The detector's shielding. The shield is comprises 3 cm archeological lead, 22 cm low activity lead, and 40 cm high-density polyethylene, encased in a stainless steel skin.

The detector was constructed and commissioned in LSM. Commissioning data was collected with the detector installed in a temporary shielding. Analysis of approximately 10 days of data collected with the detector filled with 135 mbar CH_4 is currently ongoing, with an update reported in Ref. [14]. Following this commissioning phase, the detector was shipped to SNOLAB, Canada. NEWS-G have recently begun collecting physics-quality data with the detector installed inside its compact shielding [14].

To support current and future DM searches, several developments have been necessary. In order to efficiently collect electrons generated in the 140 cm detector and improve detector

stability, a multi-anode read-out sensor, ACHINOS, has been developed [15–17]. Gas purification methods to remove contaminant gases, such as oxygen, water and radon, have also been studied, as have the production of purifiers with reduced radon emanation [18, 19]. Furthermore, several measurements and estimations of the ionisation quenching factor have recently been performed, using various methods [20–22]. Further campaigns to extend measurements to even lower energies and perform measurements in other relevant gases are planned in the near future.

2 Future fully electroformed detectors

The current NEWS-G detector's dominant background is the radioactivity contamination of the 99.99% pure copper used for the vessel, which is either inherent to the material or cosmogenically induced during its transportation [12]. The next generation NEWS-G detectors will be constructed using ultra-pure copper electroforming [23], with the detector being electroformed intact - removing the need for welding, which may introduce additional background - and directly in the underground lab where it will be hosted, to minimise cosmogenic activation. This is the motivation behind the ECuME project, which will deliver a 140 cm in diameter detector, to be installed into the current detector's shielding upon completion of its physics exploitation stage. A 30 cm in diameter prototype detector will begin construction late 2022, and will define the electroforming parameters for ECuME.

A future 3 m in diameter fully electroformed detector, DarkSPHERE, is currently being designed. DarkSPHERE will employ a water-based shielding, to suppress background contributions from the shielding material. The Boulby Underground Laboratory, UK, is being targeted as a host, with the available space in the Large Experiment Cavern sufficient to house $3 \times 3 \times 3 \text{ m}^3$ shield, which would enable DarkSPHERE to achieve a background of less than 0.01 dru below 1 keV, as estimated by Geant4 [24] simulations.

The projected sensitivity of ECuME and DarkSPHERE in a DM-nucleon spin-independent cross-section versus DM mass is shown in Figure 3. A publication is under preparation which will detail the wider physics potential of DarkSPHERE, including spin-dependent nucleon interactions, DM-electron interactions [25], and other rare-event searches.

3 Summary

NEWS-G are using spherical proportional counters to shed light on unexplored DM candidates in the 0.05–10 GeV DM mass region. The current NEWS-G detector is yielding first results, and is currently taking data for its first physics campaign in SNOLAB. In parallel, the collaboration is working towards its future detectors, which will employ intact, underground electroformed detectors to suppress radioactive contamination in the detector material. A prototype for this, miniECuME, will begin construction in late 2022, with the full-scale detector ECuME, and the planned larger-scale detector, DarkSPHERE, following in the future.

Acknowledgements

Funding information This project has received support from UK Research and Innovation Science and Technology Facilities Council through grants No. ST/S000860/1, ST/V006339/1, and ST/W005611/1.




Figure 3: Projected sensitivity of current [10] and future NEWS-G detectors in DM-nucleon spin-independent cross section versus DM mass compared to current results from CDMSLite [26], CRESST-III [27], DarkSide-50 [28], and Xenon 1T (Migdal) [29].

References

- [1] J. Billard et al., *Direct detection of dark matter—APPEC committee report**, Rep. Prog. Phys. **85**, 056201 (2022), doi:[10.1088/1361-6633/ac5754](https://doi.org/10.1088/1361-6633/ac5754).
- [2] J. A. Evans, S. Gori and J. Shelton, *Looking for the WIMP next door*, J. High Energy Phys. **02**, 100 (2018), doi:[10.1007/JHEP02\(2018\)100](https://doi.org/10.1007/JHEP02(2018)100).
- [3] K. Petraki and R. R. Volkas, *Review of asymmetric dark matter*, Int. J. Mod. Phys. A **28**, 1330028 (2013), doi:[10.1142/S0217751X13300287](https://doi.org/10.1142/S0217751X13300287).
- [4] K. M. Zurek, *Asymmetric dark matter: Theories, signatures, and constraints*, Phys. Rep. **537**, 91 (2014), doi:[10.1016/j.physrep.2013.12.001](https://doi.org/10.1016/j.physrep.2013.12.001).
- [5] M. Ibe, W. Nakano, Y. Shoji and K. Suzuki, *Migdal effect in dark matter direct detection experiments*, J. High Energy Phys. **03**, 194 (2018), doi:[10.1007/JHEP03\(2018\)194](https://doi.org/10.1007/JHEP03(2018)194).
- [6] I. Katsioulas, P. Knights, J. Matthews, T. Neep, K. Nikolopoulos, R. Owen and R. Ward, *Development of a simulation framework for spherical proportional counters*, J. Instrum. **15**, C06013 (2020), doi:[10.1088/1748-0221/15/06/C06013](https://doi.org/10.1088/1748-0221/15/06/C06013).
- [7] I. Giomataris et al., *A novel large-volume spherical detector with proportional amplification read-out*, J. Instrum. **3**, P09007 (2008), doi:[10.1088/1748-0221/3/09/P09007](https://doi.org/10.1088/1748-0221/3/09/P09007).
- [8] Q. Arnaud et al., *Precision laser-based measurements of the single electron response of spherical proportional counters for the NEWS-G light dark matter search experiment*, Phys. Rev. D **99**, 102003 (2019), doi:[10.1103/PhysRevD.99.102003](https://doi.org/10.1103/PhysRevD.99.102003).

- [9] I. Katsioulas et al., *A sparkless resistive glass correction electrode for the spherical proportional counter*, J. Instrum. **13**, P11006 (2018), doi:[10.1088/1748-0221/13/11/P11006](https://doi.org/10.1088/1748-0221/13/11/P11006).
- [10] Q. Arnaud et al., *First results from the NEWS-G direct dark matter search experiment at the LSM*, Astropart. Phys. **97**, 54 (2018), doi:[10.1016/j.astropartphys.2017.10.009](https://doi.org/10.1016/j.astropartphys.2017.10.009).
- [11] Q. Arnaud et al., *Solar Kaluza-Klein axion search with NEWS-G*, Phys. Rev. D **105**, 012002 (2022), doi:[10.1103/PhysRevD.105.012002](https://doi.org/10.1103/PhysRevD.105.012002).
- [12] L. Balogh et al., *The NEWS-G detector at SNOLAB*, J. Instrum. **18**, T02005 (2023), doi:[10.1088/1748-0221/18/02/T02005](https://doi.org/10.1088/1748-0221/18/02/T02005).
- [13] L. Balogh et al., *Copper electroplating for background suppression in the NEWS-G experiment*, Nucl. Instrum. Methods Phys. Res. A: Accel. Spectrom. Detect. Assoc. Equip. **988**, 164844 (2021), doi:[10.1016/j.nima.2020.164844](https://doi.org/10.1016/j.nima.2020.164844).
- [14] F. A. Vazquez de Sola Fernandez, *Search for light wimp recoils on methane with NEWS-G, in 14th International Conference on Identification of Dark Matter*, Vienna, Austria (2022).
- [15] A. Giganon et al., *A multiball read-out for the spherical proportional counter*, J. Instrum. **12**, P12031 (2017), doi:[10.1088/1748-0221/12/12/P12031](https://doi.org/10.1088/1748-0221/12/12/P12031).
- [16] I. Giomataris et al., *A resistive ACHINOS multi-anode structure with DLC coating for spherical proportional counters*, J. Instrum. **15**, P11023 (2020), doi:[10.1088/1748-0221/15/11/P11023](https://doi.org/10.1088/1748-0221/15/11/P11023).
- [17] I. Katsioulas et al., *ACHINOS: A multi-anode read-out for position reconstruction and tracking with spherical proportional counters*, J. Instrum. **17**, C08025 (2022), doi:[10.1088/1748-0221/17/08/C08025](https://doi.org/10.1088/1748-0221/17/08/C08025).
- [18] P. O'Brien, *Optimization of processing parameters and development of a radon trapping system for the news-g dark matter detector*, MSc. Thesis, University of Alberta, Canada (2021), doi:[10.7939/r3-qr9t-9n98](https://doi.org/10.7939/r3-qr9t-9n98).
- [19] K. Altenmuller et al., *Purification efficiency and radon emanation of gas purifiers used with pure and binary gas mixtures for gaseous dark matter detectors*, in *2021 IEEE Nuclear Science Symposium and Medical Imaging Conference (NSS/MIC)*, doi:[10.1109/NSS/MIC44867.2021.9875870](https://doi.org/10.1109/NSS/MIC44867.2021.9875870).
- [20] N.-G. Collaboration et al., *Measurements of the ionization efficiency of protons in methane*, Eur. Phys. J. C **82**, 1114 (2022), doi:[10.1140/epjc/s10052-022-11063-9](https://doi.org/10.1140/epjc/s10052-022-11063-9).
- [21] L. Balogh et al., *Quenching factor measurements of neon nuclei in neon gas*, Phys. Rev. D **105**, 052004 (2022), doi:[10.1103/PhysRevD.105.052004](https://doi.org/10.1103/PhysRevD.105.052004).
- [22] I. Katsioulas, P. Knights and K. Nikolopoulos, *Ionisation quenching factors from W-values in pure gases for rare event searches*, Astropart. Phys. **141**, 102707 (2022), doi:[10.1016/j.astropartphys.2022.102707](https://doi.org/10.1016/j.astropartphys.2022.102707).
- [23] E. W. Hoppe et al., *Reduction of radioactive backgrounds in electroformed copper for ultra-sensitive radiation detectors*, Nucl. Instrum. Methods Phys. Res. A: Accel. Spectrom. Detect. Assoc. Equip. **764**, 116 (2014), doi:[10.1016/j.nima.2014.06.082](https://doi.org/10.1016/j.nima.2014.06.082).
- [24] S. Agostinelli et al., *Geant4—a simulation toolkit*, Nucl. Instrum. Methods Phys. Res. A: Accel. Spectrom. Detect. Assoc. Equip. **506**, 250 (2003), doi:[10.1016/S0168-9002\(03\)01368-8](https://doi.org/10.1016/S0168-9002(03)01368-8).

- [25] L. Hamaide and C. McCabe, *Fueling the search for light dark matter-electron scattering with spherical proportional counters*, Phys. Rev. D **107**, 063002 (2023), doi:[10.1103/PhysRevD.107.063002](https://doi.org/10.1103/PhysRevD.107.063002).
- [26] R. Agnese et al., *Search for low-mass dark matter with CDMSlite using a profile likelihood fit*, Phys. Rev. D **99**, 062001 (2019), doi:[10.1103/PhysRevD.99.062001](https://doi.org/10.1103/PhysRevD.99.062001).
- [27] A. H. Abdelhameed et al., *First results from the CRESST-III low-mass dark matter program*, Phys. Rev. D **100**, 102002 (2019), doi:[10.1103/PhysRevD.100.102002](https://doi.org/10.1103/PhysRevD.100.102002).
- [28] P. Agnes et al., *Low-mass dark matter search with the DarkSide-50 experiment*, Phys. Rev. Lett. **121**, 081307 (2018), doi:[10.1103/PhysRevLett.121.081307](https://doi.org/10.1103/PhysRevLett.121.081307).
- [29] E. Aprile et al., *Search for light dark matter interactions enhanced by the Migdal effect or Bremsstrahlung in XENON1T*, Phys. Rev. Lett. **123**, 241803 (2019), doi:[10.1103/PhysRevLett.123.241803](https://doi.org/10.1103/PhysRevLett.123.241803).