

# VERITAS highlights of observations and results

Sonal R. Patel\* for the VERITAS collaboration

Deutsches Elektronen-Synchrotron DESY, Platanenallee 6, D-15738 Zeuthen, Germany

\* [sonal.patel@desy.de](mailto:sonal.patel@desy.de)



21st International Symposium on Very High Energy Cosmic Ray Interactions

(ISVHECRI 2022)

Online, 23-28 May 2022

doi:[10.21468/SciPostPhysProc.13](https://doi.org/10.21468/SciPostPhysProc.13)

## Abstract

Located in southern Arizona, VERITAS is amongst the most sensitive detectors for astrophysical very high energy (VHE;  $E > 100$  GeV) gamma rays and has been operational since April 2007. We highlight some recent results from VERITAS observations. These include the long-term observations of the gamma-ray binaries HESS J0632+057 and LS I +61° 303, the observations of the Galactic Center region, and of the supernova remnant Cas A. We discuss the results from a decade of multi-wavelength observations of the blazar 1ES 1215+303, the EHT 2017 campaign on the M87 galaxy, the discovery of 3C 264 in VHE, and the observation of three flaring quasars. Brief highlights of the indirect dark matter searches and targets-of-opportunity (ToO) observations are also discussed. The ToO observations allow for rapid follow-up of multi-messenger alerts and astrophysical transients.



Copyright S. R. Patel.

This work is licensed under the Creative Commons

[Attribution 4.0 International License](https://creativecommons.org/licenses/by/4.0/).

Published by the SciPost Foundation.

Received 25-08-2022

Accepted 22-06-2023

Published 29-09-2023

doi:[10.21468/SciPostPhysProc.13.035](https://doi.org/10.21468/SciPostPhysProc.13.035)



Check for updates

## 1 Introduction

VERITAS is one of the most sensitive ground-based  $\gamma$ -ray observatories. It is located at the Fred Lawrence Whipple Observatory in southern Arizona (31° 40'N, 110° 57'W, 1.3 km a.s.l.). The VERITAS array has four 12-m diameter, 12-m focal length imaging atmospheric Cherenkov telescopes. Each telescope has a Davies-Cotton-design segmented mirror dish of 345 facets and focuses the Cherenkov light from particle showers onto a pixelated camera having 499 PMTs and a total field of view of 3.5°. VERITAS began full operation in April 2007. Since then, it has undergone two major upgrades. In the first upgrade, during summer 2009, one of the four telescopes was relocated to a different position [1]. In the second upgrade, in summer 2012, PMTs (with the higher quantum efficiency and shorter time profile), and a new topological trigger system were installed [2]. The current configuration of the array can detect an object having 1% of the Crab Nebula flux in  $\sim 25$  hours with a gamma-ray-photon energy resolution




Figure 1: Typical observing plan of VERITAS.

of 15-25%. For a 1 TeV photon, the 68% containment radius is  $\leq 0.1^\circ$ , with a pointing accuracy of  $< 50''$ . The details of the evolution of the performance of the VERITAS instrument with time are discussed in [3]. In order to account for the changing optical throughput and detector performance over time, signal calibration methods have recently been implemented to produce a fine-tuned instrument response functions [4].

## 2 VERITAS observing program

The VERITAS observing season spans from September to July each year. Each year, the array collects  $\sim 950$  h of good-weather data during dark time and  $\sim 250$  h of data during bright moon (illumination of 30-65%). The typical annual VERITAS observation plan breakdown is shown in Figure 1.

## 3 Galactic science highlights

**$\gamma$ -ray binaries – HESS J0632+057 and LS I +61° 303:** Based on the composition and energy output, gamma-ray binaries can be defined as systems consisting of a compact object orbiting a star (O or Be), with periodic release of large amounts of non-thermal emission at energies  $> 1$  MeV [5]. The VERITAS archive includes one of the largest data sets for the sources of this class, which includes  $\sim 260$  h and  $\sim 174$  h of good quality data (after applying weather-based time cuts) for HESS J0632+057 and LS I +61° 303, respectively.

Located at a distance of 1.1-1.7 kpc, HESS J0632+057 consists of an unknown compact object orbiting a Be star (MWC 148) with a period of  $321 \pm 5$  days [6]. The data from three major atmospheric Cherenkov telescopes: H.E.S.S. [7], MAGIC [8], and VERITAS [3] were used to study the very high energy (VHE)  $\gamma$ -ray emission from this source, resulting in a total of 450 h over 15 years, between 2004 and 2019. The VHE  $\gamma$ -ray fluxes were found to be modulated with the orbital period of  $316.7 \pm 4.4(\text{stat}) \pm 2.5(\text{sys})$ , consistent with the value




Figure 2: Orbital phase-binned light curve of LS I +61° 303. The starting MJD and the orbital period used are 43366.8 days and 26.5 days, respectively.

obtained at X-ray energies [9]. This large data set includes dense observational coverage for several orbits, which reveal short-timescale and orbit-to-orbit variability.

The  $\gamma$ -ray binary LS I +61° 303 consists of a rapidly rotating BeVe star located at  $2.65 \pm 0.09$  kpc [10] and a compact object orbiting with a period of 26.5 days [11]. The recent detection of radio pulsations from the direction of this source by the the Five-hundred-meter Aperture Spherical radio Telescope suggests that the compact object is a rotating neutron star [12]. The orbital phase light curve of nightly-binned flux points from the analysis of  $\sim 174$  h of VERITAS data above 300 GeV is shown in Figure 2. The box shown in red includes the highest state ( $\sim 30\%$  Crab Nebula flux above 300 GeV) of LS I +61° 303 occurred in October 2014, during which the flux variability on night-timescale was observed [13]. Other than this state, nearly a factor of two flux difference in the orbital phase bin of (0.55-0.65) suggests the orbit-to-orbit variability. LS I +61° 303 is detected above  $5\sigma$  in most of the bins, except phase bins (0.1-0.2), (0.2-0.3), and (0.9-1.0), where the significance is about  $4\sigma$  [14].

**Galactic center region:** The Galactic Center (GC) region hosts numerous powerful sources and potential sites of particle acceleration, including the supermassive black hole Sagittarius A\* (Sgr A\*), supernova remnants (SNRs), and pulsar wind nebulae. VERITAS observes the GC at large zenith angles ( $\geq 60^\circ$ ), resulting in an increased effective area of about four times greater than the effective area at a zenith angle of  $20^\circ$  for energies above 10 TeV, and increased in the systematic. This also raises the energy threshold for the GC analysis to about 2 TeV. 125 h of VERITAS data resulted in a detection of Sgr A\* at a significance of  $38\sigma$ . The differential spectrum is best fitted with a power law with an exponential cutoff at  $10.0^{+4.0}_{-2.0}$  TeV and a spectral index of  $2.12^{+0.22}_{-0.17}$ . The analysis of the diffuse GC ridge shows a power law spectrum extending to the highest observed energy of  $\sim 40$  TeV and a hard spectral index of  $2.19 \pm 0.20$ . This supports the evidence for a PeVatron in the GC region. A more detailed discussion of this analysis can be found in [15].

**Supernova remnant – Cassiopeia A:** SNRs are thought to be the most favorable sites for the acceleration of Galactic cosmic-rays up to PeV energies. Among the SNRs, the young SNR are considered the best candidates to be detected at VHE, since the highest energy cosmic rays,

which are believed to be produced early on, will not yet have escaped from the production site. The young ( $\sim 350$  years old) core-collapse SNR Cassiopeia A is a potential PeVatron candidate. VERITAS has studied this source with a deep exposure of 65 h. The joint spectrum was produced with VERITAS data covering 200 GeV-10 TeV and 10.8 years of *Fermi*-Large Area Telescope data covering 0.1-500 GeV. The best fit was obtained with the power law spectral index of  $2.17 \pm 0.02(\text{stat})$  and cutoff energy of  $2.3 \pm 0.5(\text{stat})$  TeV [16]. Considering a one-zone model, proton acceleration up to at least 6 TeV is required to reproduce the observed  $\gamma$ -ray spectrum.

## 4 Extra-galactic science highlights

The VERITAS AGN observations are broadly conducted under four major programs, namely: discovery program, multi-wavelength (MWL) campaigns, ToO, and cosmology. The highlights of the first three programs are mentioned in this section.

**Blazar – 1ES 1215+303:** A high-synchrotron-peaked BL Lac (HBL) 1ES 1215+303 ( $z=0.13$ , [17]) was extensively studied in a MWL context using long-term data from radio to  $\gamma$ -ray energies. a VERITAS exposure of 175.8 h was used in this study, which includes regular monitoring observations since December 2008. The synchrotron peak frequency from a low state to the 2017 flaring state was observed to be shifted from infrared to soft X-ray [18].

**Radio Galaxies – 3C 264 and M87:** The misaligned geometry of radio galaxies provides a unique view of the AGN's jet and super massive back hole. Among TeV-detected radio galaxies, 3C 264 ( $z = 0.0217$ , [19]) is the most distant, and is only the fourth known radio galaxy at these energies. VERITAS has collected  $\sim 57$  h good quality data between February 2017 and May 2019, which yielded a detection with a statistical significance of  $7.8\sigma$ . The VHE  $\gamma$ -ray spectrum was well described by a power law with an index of  $2.20 \pm 0.27$  and a flux of  $\sim 0.7\%$  of the Crab Nebula flux above 315 GeV [20].

During the 2017 Event Horizon Telescope MWL campaign on M87, VERITAS captured the source in its historically low state, but still dominating over the nearest knot, HST-1 [21]. The most complete simultaneous MWL spectrum was reported in that study, which included 15 h of quality-selected VERITAS data. The analysis resulted in an overall statistical significance of  $3.8\sigma$ . The legacy data set and analysis scripts were made available to the community through the Cyverse repository [22].

**Quasars – 3C 279, PKS 1222+216, and Ton 599:** Known flat-spectrum radio quasars (FSRQs) are generally at larger distances compared to BL Lacs, hence with the current generation VHE  $\gamma$ -rays telescopes their detection is difficult, both because of the attenuation of VHE  $\gamma$ -rays from these distances due to absorption by the extra-galactic background light and because of possible intrinsic spectral cutoffs above GeV energies [23]. Three FSRQs; 3C 279, PKS 1222+216, and Ton 599, were studied to explore the  $\gamma$ -ray variability and spectral characteristics using almost 100 h of VERITAS data spanning over 10 years. The location of the  $\gamma$ -ray emission region and the jet Doppler factor were constrained during VHE-detected flares in 2014 and 2017, for PKS 1222+216 and Ton 599, respectively [24]. Also, theoretical constraints on the potential production of PeV-scale neutrinos were placed during these VHE flares.




Figure 3: The sky map showing some MM triggers observed by VERITAS (Adopted from [27]).

## 5 Indirect dark matter search

Dwarf Spheroidal galaxies (dSphs) are a favorable target class for an indirect dark matter (DM) search. The analysis of four dSphs (Boötes I, Draco, Segue 1, and Ursa Minor) with the VERITAS data taken from 2007 to 2013 was reported in [25]. This included a total quality-selected observation time of 476 h to search for a DM signal and was sensitive to potential signals in the  $\tau^+\tau^-$  and  $b\bar{b}$  annihilation channels. No DM signal was detected.

## 6 Multi-messenger and astrophysical transient observations

VERITAS devotes a part of its observing time to multi-messenger (MM) and astrophysical transient observations to search for electromagnetic counterparts to high energy neutrinos and gravitational waves. These observations are proposal driven and result into significant fraction of total VERITAS observing time. Figure 3 shows some MM triggers observed by VERITAS. The IceCube observatory reported a well-reconstructed high energy neutrino event, IceCube-201114A (GCN 28887), on November 14, 2020, having an estimated energy of  $\sim 214$  TeV. It is spatially coincident with the high-energy-peaked object, NVSS J065844+063711 [26]. During November 15-19, 2020, VERITAS observed NVSS J065844+063711, and a differential upper limit was reported in [26] using 7 h quality-selected data.

## 7 Summary and Conclusions

VERITAS has a strong and multi-faceted science program in the  $\gamma$ -ray band. Selected science results are highlighted, covering results from Galactic, extra-galactic, fundamental physics, and multi-messenger observations. VERITAS is operating extremely well and continues to provide high quality VHE  $\gamma$ -ray data and scientific results to the community. VERITAS has been recommended for the next cycle of NSF operations funding through 2025.

## Acknowledgments

We acknowledge the excellent work of the technical support staff at the Fred Lawrence Whipple Observatory and at the collaborating institutions in the construction and operation of the instrument.

**Funding information** This research is supported by grants from the U.S. Department of Energy Office of Science, the U.S. National Science Foundation and the Smithsonian Institution, by NSERC in Canada, and by the Helmholtz Association in Germany. This research used resources provided by the Open Science Grid, which is supported by the National Science Foundation and the U.S. Department of Energy's Office of Science, and resources of the National Energy Research Scientific Computing Center (NERSC), a U.S. Department of Energy Office of Science User Facility operated under Contract No. DE-AC02-05CH11231.

## References

- [1] J. S. Perkins et al., *VERITAS telescope 1 relocation: Details and improvements*, (arXiv preprint) doi:[10.48550/arXiv.0912.3841](https://doi.org/10.48550/arXiv.0912.3841).
- [2] B. Zitzer, *The VERITAS upgraded telescope-level trigger systems: Technical details and performance characterization*, (arXiv preprint) doi:[10.48550/arXiv.1307.8360](https://doi.org/10.48550/arXiv.1307.8360).
- [3] N. Park, *Performance of the VERITAS experiment*, Proc. Sci. **236**, 771 (2016), doi:[10.22323/1.236.0771](https://doi.org/10.22323/1.236.0771).
- [4] C. B. Adams et al., *The throughput calibration of the VERITAS telescopes*, Astron. Astrophys. **658**, A83 (2022), doi:[10.1051/0004-6361/202142275](https://doi.org/10.1051/0004-6361/202142275).
- [5] G. Dubus, *Gamma-ray binaries and related systems*, Astron. Astrophys. Rev. **21**, 64 (2013), doi:[10.1007/s00159-013-0064-5](https://doi.org/10.1007/s00159-013-0064-5).
- [6] S. D. Bongiorno et al., *A new TeV binary: The discovery of an orbital period in HESS J0632+057*, Astrophys. J. **737**, L11 (2011), doi:[10.1088/2041-8205/737/1/L11](https://doi.org/10.1088/2041-8205/737/1/L11).
- [7] J. A. Hinton, *The status of the HESS project*, New Astron. Rev. **48**, 331 (2004), doi:[10.1016/j.newar.2003.12.004](https://doi.org/10.1016/j.newar.2003.12.004).
- [8] J. Aleksić et al., *The major upgrade of the MAGIC telescopes, part II: A performance study using observations of the Crab nebula*, Astropart. Phys. **72**, 76 (2016), doi:[10.1016/j.astropartphys.2015.02.005](https://doi.org/10.1016/j.astropartphys.2015.02.005).
- [9] C. B. Adams et al., *Observation of the gamma-ray binary HESS J0632 + 057 with the H.E.S.S., MAGIC, and VERITAS telescopes*, Astrophys. J. **923**, 241 (2021), doi:[10.3847/1538-4357/ac29b7](https://doi.org/10.3847/1538-4357/ac29b7).
- [10] L. Lindegren et al., *Gaia early data release 3*, Astron. Astrophys. **649**, A2 (2021), doi:[10.1051/0004-6361/202039709](https://doi.org/10.1051/0004-6361/202039709).
- [11] J. Casares, I. Ribas, J. M. Paredes, J. Martí and C. A. Prieto, *Orbital parameters of the microquasar LS I + 61 303*, Mon. Not. R. Astron. Soc. **360**, 1105 (2005), doi:[10.1111/j.1365-2966.2005.09106.x](https://doi.org/10.1111/j.1365-2966.2005.09106.x).

- [12] S.-S. Weng et al., *Radio pulsations from a neutron star within the gamma-ray binary LS I + 61° 303*, Nat. Astron. **6**, 698 (2022), doi:[10.1038/s41550-022-01630-1](https://doi.org/10.1038/s41550-022-01630-1).
- [13] S. Archambault et al., *Exceptionally bright TeV flares from the binary LS I + 61° 303*, Astrophys. J. **817**, L7 (2016), doi:[10.3847/2041-8205/817/1/L7](https://doi.org/10.3847/2041-8205/817/1/L7).
- [14] D. Kieda et al., *Very high energy gamma-ray emission from the binary system LS I + 61 303*, Proc. Sci. **395**, 832 (2021), doi:[10.22323/1.395.0832](https://doi.org/10.22323/1.395.0832).
- [15] C. B. Adams et al., *VERITAS observations of the Galactic Center region at multi-TeV gamma-ray energies*, Astrophys. J. **913**, 115 (2021), doi:[10.3847/1538-4357/abf926](https://doi.org/10.3847/1538-4357/abf926).
- [16] A. U. Abeysekara et al., *Evidence for proton acceleration up to TeV energies based on VERITAS and Fermi-LAT observations of the Cas A SNR*, Astrophys. J. **894**, 51 (2020), doi:[10.3847/1538-4357/ab8310](https://doi.org/10.3847/1538-4357/ab8310).
- [17] A. E. Truebenbach and J. Darling, *The VLBA extragalactic proper motion catalog and a measurement of the secular aberration drift*, Astrophys. J. Suppl. **233**, 3 (2017), doi:[10.3847/1538-4365/aa9026](https://doi.org/10.3847/1538-4365/aa9026).
- [18] J. Valverde et al., *A decade of multiwavelength observations of the TeV blazar 1ES 1215 + 303: Extreme shift of the synchrotron peak frequency and long-term optical-gamma-ray flux increase*, Astrophys. J. **891**, 170 (2020), doi:[10.3847/1538-4357/ab765d](https://doi.org/10.3847/1538-4357/ab765d).
- [19] R. J. Smith, J. R. Lucey, M. J. Hudson, D. J. Schlegel and R. L. Davies, *Streaming motions of galaxy clusters within 12 000 km s<sup>-1</sup> – I. New spectroscopic data*, Mon. Not. R. Astron. Soc. **313**, 469 (2000), doi:[10.1046/j.1365-8711.2000.03251.x](https://doi.org/10.1046/j.1365-8711.2000.03251.x).
- [20] A. Archer et al., *VERITAS discovery of VHE emission from the radio galaxy 3C 264: A multiwavelength study*, Astrophys. J. **896**, 41 (2020), doi:[10.3847/1538-4357/ab910e](https://doi.org/10.3847/1538-4357/ab910e).
- [21] J. C. Algaba et al., *Broadband multi-wavelength properties of M87 during the 2017 event horizon telescope campaign*, Astrophys. J. Lett. **911**, L11 (2021), doi:[10.3847/2041-8213/abef71](https://doi.org/10.3847/2041-8213/abef71).
- [22] EHT MWL Science Working Group, *Multi-wavelength observations of M87 during the 2017 event horizon telescope campaign*, CyVerse Data Commons (2021), doi:[10.25739/mhh2-cw46](https://doi.org/10.25739/mhh2-cw46).
- [23] B. E. Stern and J. Poutanen, *The mystery of spectral breaks: Lyman continuum absorption by photon-photon pair production in the Fermi GeV spectra of bright blazars*, Astrophys. J. **794**, 8 (2014), doi:[10.1088/0004-637X/794/1/8](https://doi.org/10.1088/0004-637X/794/1/8).
- [24] C. B. Adams et al., *Variability and spectral characteristics of three flaring gamma-ray quasars observed by VERITAS and Fermi-LAT*, Astrophys. J. **924**, 95 (2022), doi:[10.3847/1538-4357/ac32bd](https://doi.org/10.3847/1538-4357/ac32bd).
- [25] C. Giuri, *VERITAS dark matter search in dwarf spheroidal galaxies: An extended source analysis*, Proc. Sci. **395**, 515 (2021), doi:[10.22323/1.395.0515](https://doi.org/10.22323/1.395.0515).
- [26] R. de Menezes et al., *Multi-messenger observations of the Fermi-LAT blazar 4FGL J0658.6 + 0636 consistent with an IceCube high-energy neutrino*, Proc. Sci. **358**, 955 (2021), doi:[10.22323/1.395.0955](https://doi.org/10.22323/1.395.0955),
- [27] M. Santander, *Recent results from the VERITAS multimessenger program*, Proc. Sci. **358**, 782 (2019), doi:[10.22323/1.358.0782](https://doi.org/10.22323/1.358.0782).